

EE5130: Detection and Estimation Theory

Problem Set 8

1. Consider the observed data set

$$Y_k = A + W_k \quad k = 1, \dots, n$$

where A is an unknown level, which is assumed to be positive ($A > 0$) and W_k is WGN with unknown variance A . Find the variance of the sample mean estimator and compare it to the CRLB. Does the sample mean estimator attain the CRLB for finite n ? How about if $n \rightarrow \infty$? Is the MLE or the sample mean a better estimator?

2. A formal definition of the consistency of an estimator is given as follows. An estimator $\hat{\theta}$ is consistent if, given any $\epsilon > 0$,

$$\lim_{n \rightarrow \infty} Pr\{|\hat{\theta} - \theta| > \epsilon\} = 0$$

Prove that the sample mean is a consistent estimator for the problem of estimating a DC level A in white Gaussian noise of known variance. Hint : Use Chebychev's inequality: For a random variable X with mean μ and variance σ^2 , $P(|X - \mu| \geq \epsilon) \leq \frac{\sigma^2}{\epsilon^2}$.

3. If we observe n IID samples from a Bernoulli experiment (coin toss) with the probabilities

$$\begin{aligned} Pr\{Y_n = 1\} &= p \\ Pr\{Y_n = 0\} &= 1 - p \end{aligned}$$

find the MLE of p .

4. For n IID observations from a $\mathcal{U}[0, \theta]$ PDF, find the MLE of θ .
5. For N IID observations from a $\mathcal{N}(0, \frac{1}{\theta})$ PDF, where $\theta > 0$, find the MLE of θ and its asymptotic PDF.
6. For N IID observations from the PDF $\mathcal{N}(A, \sigma^2)$, where A and σ^2 are both unknown, find the MLE of the SNR $\alpha = \frac{A^2}{\sigma^2}$.